

# Diverse Host Feeding on Nesting Birds May Limit Early-Season West Nile Virus Amplification

Andrea M. Egizi,<sup>1,2</sup> Ary Farajollahi,<sup>1,3</sup> and Dina M. Fonseca<sup>1,2</sup>

## Abstract

Arboviral activity tracks vector availability, which in temperate regions means that transmission ceases during the winter and must be restarted each spring. In the northeastern United States, *Culex restuans* Theobald resumes its activity earlier than *Culex pipiens* L. and is thought to be important in restarting West Nile virus (WNV) transmission. Its role in WNV amplification, however, is unclear, because viral levels commonly remain low until the rise of *Cx. pipiens* later in the season. Because a vector's feeding habits can reveal key information about disease transmission, we identified early-season (April–June) blood meals from *Cx. restuans* collected throughout New Jersey, and compared them to published datasets from later in the season and also from other parts of the country. We found significantly higher avian diversity, including poor WNV hosts, and fewer blood meals derived from American Robins (17% versus over 40% found in later season). Critically, we identified blood meals from significantly more female than male birds in species where females are the incubating sex, suggesting that *Cx. restuans* is able to feed on such a wide variety of hosts in early spring because incubating birds are easy targets. Because WNV amplification depends on virus consistently reaching competent hosts, our results indicate that *Cx. restuans* is unlikely to be an amplifying vector of WNV in the early season. As the season progresses, however, changes in the availability of nesting birds may make it just as capable as *Cx. pipiens*, although at somewhat lower abundance as the summer progresses.

**Key Words:** Blood meal analysis—West Nile virus—Dilution effect—Arbovirus—Vector ecology.

## Introduction

IT IS WELL ESTABLISHED THAT West Nile virus (WNV), like St. Louis encephalitis virus (SLE), is overwhelmingly associated with *Culex* mosquitoes (Campbell et al. 2002, Nasci et al. 2002), and three species (*Culex pipiens* L., *Culex quinquefasciatus* Say, and *Culex tarsalis* Coquillett) have been incriminated as the primary vectors in various regions. A fourth species, *Culex restuans* Theobald, is abundant in many areas in the northeastern United States and has been shown to be a competent WNV vector (Ebel et al. 2005). *Cx. restuans* starts its activity earlier in the year than *Cx. pipiens*, and studies in different regions of the United States reported temporally distinct peaks of abundance for these two species (Lee and Rowley 2000, Andreadis et al. 2001, Lampman et al. 2006). In the eastern-central United States, the cross-over date typically occurs in early July, with *Cx. restuans* populations being more abundant in May and June and

*Cx. pipiens* in July and August (Andreadis et al. 2001, Gong et al. 2011).

In climates with cold winters that suppress vector activity, the cycle of transmission of arboviruses like WNV and SLE needs to be reinitiated each year. Because *Cx. restuans* are among the first bird biters to emerge from diapause in the northeastern United States, they are considered critical for initiating WNV transmission among birds in early spring, whereas *Cx. pipiens* amplifies viral levels later in the season (Andreadis et al. 2001). The same scenario has been proposed by Reiter (1988) for the relative roles of *Cx. restuans* and *Cx. pipiens* in the transmission of SLE. Although extrinsic factors such as temperature and rainfall affect the intensity of WNV transmission (Kramer et al. 2008), it is unclear why WNV levels remain low until summer, even after temperatures permissible for viral replication have been reached (Dohm et al. 2002, Kilpatrick et al. 2008). One way to investigate the early-season amplification of WNV is to examine the composition and diversity of *Cx. restuans* blood

<sup>1</sup>Center for Vector Biology and <sup>2</sup>Graduate Program in Ecology and Evolution, Department of Entomology, Rutgers University, Brunswick, New Jersey.

<sup>3</sup>Mercer County Mosquito Control, West Trenton, New Jersey.

meal hosts during this time. Most passerine species are competent WNV hosts, whereas many other groups of birds, such as doves and shorebirds, are not (Komar et al. 2003, Blitvich 2008), and higher bird species diversity has been linked to lower WNV transmission (Ezenwa et al. 2006, Allan et al. 2009), a phenomenon known as the dilution effect (Ostfeld and Keesing 2000).

In this study, we examined the blood-feeding patterns of *Cx. restuans* collected in the early season in locations throughout New Jersey over a 10-year period. We analyzed the species diversity and sex ratio of blood meal hosts and characteristics of individual bird species, such as WNV competence, presence of migratory behavior, and egg incubation. We also compared our results to blood-feeding studies of *Cx. restuans* collected later in the season from the same region, as well as in Illinois and Tennessee. The latter locations were an effort to account for differences in bird assemblages and identify temporal patterns.

## Methods

### Collection of mosquitoes

Due to the difficulty of obtaining blooded mosquitoes in the early season, the specimens used in this analysis were collected by collaborators from a variety of locations throughout 12 New Jersey counties over 10 years, 2001–2011. Specimens were mostly collected as part of local surveillance efforts by mosquito control agencies using light traps, gravid traps, resting boxes, BioGents Sentinel traps (Biogents, AG, Regensburg, Germany), and backpack aspiration. All blooded specimens identified as *Cx. restuans* on the basis of morphology were immediately placed in a  $-80^{\circ}\text{C}$  freezer and held for extraction and molecular identification. A subset collected between April 15 and June 15 across all years was analyzed in this study ( $n=113$ ). We chose to not identify the blood meals of late-season *Cx. restuans* because specimens collected in the same manner and from many of the same sites as ours have already been analyzed and published (Apperson et al. 2004) and are included in the late season “Northeast” dataset (see below).

### Identification of blood meals

Blooded abdomens were removed over ice, and DNA was extracted using a Qiagen DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA). Only specimens confirmed to be *Cx. restuans* (105/113 specimens) using a molecular assay (Crabtree et al. 1995) were included in the final analysis. Specimens were first screened with BM primers (Kocher et al. 1989), and those that did not amplify were subsequently tested with Avian A (Cicero and Johnson 2001), then Rep/Amph (Cupp et al. 2004), and finally Mammal A primers (Ngo and Kramer 2003) following established protocols. Amplification products were sequenced with the forward primer of each pair. We identified the putative source of the blood using a GenBank BLAST search with a match  $\geq 98\%$  (Kent 2009). Mixed sequences were cloned and sent for sequencing. A total of 83 specimens (79.0%) were successfully identified using these methods.

To identify additional samples, a smaller 130-bp fragment was amplified using the primers CytbVertR1 (Egizi et al. 2013) and BMF1 (originally called H15149; Kocher et al.

1989). One mixed sequence amplified with these primers was cloned. This approach allowed the additional identification of 11 blood meals for a total of 94 (89.5%) identified blood meals in *Cx. restuans*.

### Comparison with previous studies

Existing data on *Cx. restuans* avian blood-feeding habits was obtained from published studies and pooled by geographic location into three datasets. The Northeast data set encompassed four studies conducted in New Jersey, New York, and Connecticut (Apperson et al. 2002, Apperson et al. 2004, Molaei et al. 2006, Diuk-Wasser et al. 2010); the IL dataset was from two studies in southwest Chicago, Illinois (Hamer et al. 2008, Hamer et al. 2009); and the TN dataset was from Shelby County, Tennessee (Savage et al. 2007). Only studies that used molecular techniques to distinguish *Cx. restuans* from *Cx. pipiens* and that published a breakdown of all avian blood meals by species were included. These datasets were compared with the early-season New Jersey data from our study (called NJearly).

We analyzed a table of counts of each bird species in each dataset using a correspondence analysis with the *ca* package (Nenadic and Greenacre 2007) in the statistical software program R (R Core Team 2013). Proportion of American Robin blood meals was compared across datasets, as robins have been implicated as an important amplification host of WNV (Kilpatrick et al. 2006a). Because the presence of migrant birds may also affect WNV transmission, we calculated the proportion of migratory bird blood meals. A species was considered migratory for a particular dataset if it is not considered a year-round resident in the area where that dataset was collected (Poole 2005). Finally, because of their ability to dampen WNV transmission, we also calculated the proportion of nonpasserine blood meals in each dataset. Each proportion (robin, migratory, and nonpasserine) in the published datasets was compared to the early-season New Jersey data in a chi-squared test.

We calculated Rényi profiles, a graphical means of comparing species diversity between samples, in R (R Core Team 2013) using the packages *BiodiversityR* (Kindt and Coe 2005) and *vegan* (Oksanen et al. 2013). Rényi profiles plot Rényi entropy ( $H_{\alpha}$ ) over a range of alphas ( $\alpha$ ) using the formula:

$$H_{\alpha} = \frac{\ln \sum_{i=1}^S p_i^{\alpha}}{1 - \alpha}$$

where  $p_i$  is the partial abundance of species  $i$  and  $S$  is the total number of species in the dataset (Rényi 1961, Tóthmérész 1995). For different values of  $\alpha$ , this measure is related to several well-known measures of diversity including:  $\alpha=0$  for simple species richness,  $\alpha=1$  for Shannon diversity,  $\alpha=2$  for the Simpson index, and  $\alpha=\infty$  for Berger–Parker diversity (Tóthmérész 1995). Therefore, Rényi profiles present a more comprehensive picture of diversity than looking at any measure individually and allow diversity to be ranked among sites (Tóthmérész 1995, Kindt and Coe 2005). Other diversity measures calculated were Shannon diversity (Shannon and Weaver 1949), inverse Simpson diversity (Simpson 1949), and the Pielou measure of evenness (Pielou 1966), using *BiodiversityR* and *vegan* (Kindt and Coe 2005, Oksanen et al. 2013).

TABLE 1. AVIAN BLOOD MEALS IDENTIFIED IN *CULEX RESTUANS* COLLECTED IN THE EARLY SEASON (APRIL 15–JUNE 15) IN NEW JERSEY, 2001–2011

Avian species	Code	No. meals	% (n = 96)
American Robin ( <i>Turdus migratorius</i> ) <sup>†a</sup>	AMRO	16	16.67
Cooper's Hawk ( <i>Accipiter cooperii</i> ) <sup>†a</sup>	COHA	11	11.46
Gray Catbird ( <i>Dumetella carolinensis</i> )	GRCA	10	10.42
Wood Thrush ( <i>Hylocichla mustelina</i> )	WOTH	6	6.25
Common Grackle ( <i>Quiscalus quiscula</i> )	COGR	6	6.25
European Starling ( <i>Sturnus vulgaris</i> )	EUST	6	6.25
House Sparrow ( <i>Passer domesticus</i> )	HOSP	5	5.21
Mourning Dove ( <i>Zenaida macroura</i> )	MODO	4	4.17
Swamp Sparrow ( <i>Melospiza georgiana</i> )	SWSP	4	4.17
Northern Cardinal ( <i>Cardinalis cardinalis</i> )	NOCA	4	4.17
House Finch ( <i>Carpodacus mexicanus</i> ) <sup>‡b</sup>	HOFI	4	4.17
Barn Swallow ( <i>Hirundo rustica</i> )	BASW	3	3.13
Tufted Titmouse ( <i>Baeolophus bicolor</i> ) <sup>†a</sup>	TUTI	3	3.13
Brown-headed Cowbird ( <i>Molothrus ater</i> )	BRCO	2	2.08
Common Yellowthroat ( <i>Geothlypis trichas</i> )	COYE	2	2.08
Great Egret ( <i>Ardea alba</i> )	GREG	2	2.08
American Crow ( <i>Corvus brachyrhynchos</i> )	AMCR	2	2.08
Red-winged Blackbird ( <i>Agelaius phoeniceus</i> )	REBL	1	1.04
Blue Grosbeak ( <i>Passerina caerulea</i> )	BLGR	1	1.04
Tennessee Warbler ( <i>Vermivora peregrina</i> )	TEWA	1	1.04
Veery ( <i>Catharus fuscescens</i> )	VEER	1	1.04
Yellow-throated Warbler ( <i>Dendroica dominica</i> )	YEWA	1	1.04
Ross's Goose ( <i>Anser rossii</i> )	ROGO	1	1.04

“Code” is the four-letter code used for that species in Fig. 1.

<sup>a</sup>Species found in three-way mixed meal.

<sup>b</sup>Species found in a mixed meal with human.

#### Blood meal sexing

We used the assay described by Griffiths et al. (1998), which relies on a size discrepancy between the CHD-Z and CHD-W alleles (male birds are ZZ and females are ZW) (Griffiths et al. 1998) and works on a large variety of bird species. However, because in Cooper's Hawks the target fragments differ by only 2 base pairs (GenBank accession nos. JX460775 and JX460774) after amplification, we digested those PCR products with DraI (New England Biolabs, Beverly, MA). DraI cuts only the CHD-W allele, producing three bands in females and one in males. All PCR products and digestion products were run on a 3% agarose gel for 90 min to ensure separation of DNA bands.

Male and female blood meals were pooled across all avian species and tested in a chi-squared test against a null hypothesis of 1:1 male:female. A second chi-squared test omitted all species with male incubation, identified using the Birds of North America Online database (Poole 2005), as well as the Brown-headed Cowbird, a nest parasite that does not incubate its own eggs.

#### Results


We identified blood meals from 94 *Cx. restuans* mosquitoes to species. Two specimens (~2%) contained mixed blood meals—one a two-way mix between human and House Finch and the other a three-way mix between American Robin, Tufted Titmouse, and Cooper's Hawk. All other mixes were cases where the primers amplified mosquito DNA instead of that of the vertebrate host, a phenomenon we have observed when the vertebrate DNA is highly degraded (A.E. Egizi, personal observation). The human blood identified in the mix with House Finch was the only mammalian meal detected in our studies from *Cx. restuans*. The remaining blood meals contained 96 bird sequences from 23 avian species (Table 1).

The two datasets in the northeast United States (NJearly and Northeast) had similarly high species richness compared to IL and TN, although sample sizes were significantly smaller (Table 2). Importantly, however, the NJearly dataset had a more even distribution of species compared to the other three resulting in the highest Shannon and Simpson diversity

TABLE 2. COMPARISON OF DIVERSITY INDICES AND OTHER CHARACTERISTICS OF *CULEX RESTUANS* BLOOD MEAL DATASETS

Dataset	N	S	H'	1/D	J'	$\hat{p}$ robin	$\hat{p}$ nonpasserine	$\hat{p}$ migratory
NJearly	96	23	2.81	12.91	0.896	0.17	0.19	0.19
Northeast	98	26	2.38	4.73	0.729	0.44	0.14	0.17
IL	211	18	1.89	3.92	0.654	0.45	0.07	0.04
TN	203	14	1.53	2.85	0.580	0.55	0.02	0.04

N, number of blood meals; S, avian species richness; H', Shannon information index; 1/D, inverse Simpson diversity index; J'Pielou's measure of evenness, calculated as  $H'/\ln(S)$ ;  $\hat{p}$ , the proportion of blood meals with a particular characteristic.



**FIG. 1.** Rényi diversity profile values ( $H_\alpha$ ) plotted over a range of  $\alpha$  for each blood meal dataset. Smaller values of  $\alpha$  tend to weigh species richness more heavily, whereas larger values weigh evenness more heavily. Shannon and Simpson diversity, which include both factors in their calculation of diversity, fall into the middle ( $\alpha=1$  and  $\alpha=2$ , respectively). Northeast = the dataset comprising five studies conducted in New York, New Jersey, and Connecticut; NJearly = the dataset from this paper; IL = dataset comprising two studies in and around Chicago, IL; TN = dataset from Shelby County, TN.

indices (Table 2). These patterns are reflected in the Rényi diversity profiles in Figure 1, where the NJearly profile has a flatter slope, indicating greater evenness than the other communities. Because the TN dataset is consistently lower across all values of  $\alpha$ , it has the lowest overall diversity, followed by IL; however, the top two profiles cannot be definitively ordered using this method because they intersect (Fig. 1).

The NJearly dataset has the highest evenness because of a significantly lower proportion of American Robins (Tables 2 and 3). Both the NJearly and northeast datasets had significantly higher proportions of migratory birds and nonpasserine birds, compared to the TN or IL data (Tables 2 and 3). In a correspondence analysis, NJearly is separated from the other three datasets along Axis 1, which makes up 50% of the variation (Fig. 2). Note that American Robin (AMRO) falls near the other datasets along Axis 1 (Fig. 2).

The sex of the blood meal host was determined in 80/94 samples (Table 4). The overall sex ratio was not significantly different from 1:1 in a chi-squared test ( $\chi^2 = 1.25$ , degrees of freedom [df] = 1,  $0.50 < p < 0.25$ ). However, the sex ratio was

**TABLE 3.** CHI-SQUARED VALUES FOR TESTS COMPARING OBSERVED PROPORTIONS ( $\hat{p}$ ) OF *Cx. RESTUANS*-DERIVED BLOOD MEALS IN NEW JERSEY EARLY SEASON DATA WITH OTHER PUBLISHED DATASETS

Dataset	$\hat{p}$ robin	$\hat{p}$ nonpasserine	$\hat{p}$ migratory
Northeast	16.97**	0.70	0.218
IL	23.67**	49.66**	23.25**
TN	39.47**	59.60**	20.14**

\*\* $p < 0.001$ .

significantly female-biased in species with female-only incubation ( $\chi^2 = 6.12$ ,  $df = 1$ ,  $p < 0.05$ ).


## Discussion

Our study shows that early-season *Cx. restuans* collected in New Jersey feed almost exclusively on birds, although we did find a single human meal present in a mix. Previous studies of *Cx. restuans* in New York (Patrican et al. 2007) and Connecticut (Molaei et al. 2006, Diuk-Wasser et al. 2010) found exclusively avian meals, whereas other studies found as much as 13–32% mammalian (Apperson et al. 2002, Apperson et al. 2004, Savage et al. 2007, Hamer et al. 2008, Hamer et al. 2009). The frequency of mixed meals we observed (2%) is within the range found in previous studies (1–4%) (Apperson et al. 2002, Savage et al. 2007, Hamer et al. 2009). The three-way mixed meal we identified is a unique find and may have been possible because our primers targeting a smaller DNA fragment (130 bp) facilitated detection of degraded DNA. Our dataset and the Northeast dataset had a significantly greater proportion of migratory birds (19% and 17%, respectively) than either the IL or TN data (4% for both). This is probably due to geography, because the mid Atlantic coast (particularly the Cape May and Delmarva peninsulas) is considered a major stopover for neotropical migrants (Mabey et al. 1993). In New Jersey, only 22 out of 350 avian species detected annually in the state are permanent residents (Dunne et al. 1989), so the majority of the 210 birds observed breeding in the state each year (Boyle 2011) are migrants.

Numerous studies from multiple parts of the country have found that *Culex* mosquitoes derive a large portion of their meals from American Robins (Apperson et al. 2002, Apperson et al. 2004, Kilpatrick et al. 2006, Molaei et al. 2006, Savage et al. 2007, Hamer et al. 2008, Hamer et al. 2009, Kent et al. 2009, Diuk-Wasser et al. 2010, Montgomery et al. 2011). They also seem to prefer robins in laboratory choice tests (Simpson et al. 2009) and feed in greater frequency than predicted by its abundance in the field, leading researchers to postulate that robins are major amplification hosts of WNV (Kilpatrick et al. 2006a). In our study of early-season *Cx. restuans*, although robins are still the most abundant species at 17%, this number is significantly lower than in the other three datasets (Tables 2 and 3).

It is possible this difference is a consequence of our broad sampling approach, collecting mosquitoes from 12 different counties and in many different types of environments. Most studies focus their efforts in a smaller area, often in urban and suburban neighborhoods where WNV has been detected. In New Jersey, residential areas contain significantly more robins and House Sparrows than adjacent wetland areas (Johnson et al. 2012), so sampling in residential areas might yield more robin blood meals. However, the Northeast dataset comprised four different studies in three different states, with a range of environments (of note, several of the specimens were obtained from the same sources as ours), and in fact had similar species richness and sample size to our study, whereas TN and IL had lower species richness despite larger sample sizes. This indicates that at least for the comparison between our data and the Northeast data, collection bias does not explain the difference in diversity patterns.

More likely, then, these differences result from divergent host behavior in the early season. Changes in the behavior of



**FIG. 2.** Correspondence analysis biplot of host species (grey triangles) and location dataset (black circles) scores in dimensions 1 and 2 explaining 49.69% and 26.74% of the variance, respectively. Species codes follow Table 1, with additional acronyms for those not in this dataset but in others as follows: BAOR, Baltimore Oriole; BLCH, Black-capped Chickadee; BLHR, Black-crowned Night Heron; BLJA, Blue Jay; BLVI, Blue-headed Vireo; BRTH, Brown Thrasher; CACH, Carolina Chickadee; CAGO, Canada Goose; CEWA, Cedar Waxwing; EABL, Eastern Bluebird; FISP, Field Sparrow; GRHE, Green Heron; MALL, Mallard; NOFL, Northern Flicker; NOMO, Northern Mockingbird; PRWA, Prairie Warbler; ROGR, Rose-breasted Grosbeak; SCTA, Scarlet Tanager; SHHA, Sharp-shinned Hawk; SOSP, Song Sparrow; WITU, Wild Turkey. Species that are strongly associated with a particular dataset are clustered near it on the plot.

**TABLE 4. CULEX RESTUANS BLOOD MEALS FOR WHICH THE SEX WAS IDENTIFIED USING A PCR ASSAY, BROKEN DOWN BY SPECIES**

Avian species	No. females identified	No. males identified	Incubating sex
American Crow	2	0	Female
American Robin	8	5	Female
Blue Grosbeak	1	0	Female
Common Grackle	3	2	Female
Common Yellowthroat	1	1	Female
Cooper’s Hawk	8	2	Female
Gray Catbird	5	4	Female
House Finch	1	1	Female
Northern Cardinal	2	0	Female
Ross’s Goose	0	1	Female
Swamp Sparrow	2	1	Female
Tennessee Warbler	1	0	Female
Tufted Titmouse	0	2	Female
Wood Thrush	5	1	Female
<b>Total female incubation</b>	<b>39</b>	<b>20</b>	
Barn Swallow	1	2	Both
European Starling	2	4	Both
Great Egret	0	2	Both
House Sparrow	1	4	Both
Mourning Dove	1	3	Both
Brown-headed Cowbird	1	0	N/A
<b>Grand total</b>	<b>45</b>	<b>35</b>	

robins at the conclusion of the breeding season have been invoked to explain why *Cx. pipiens* shift their feeding away from robins in late summer (Kilpatrick et al. 2006b, Diuk-Wasser et al. 2010). Conversely, it is possible that the early-season behavior of robins makes them less available as hosts, or more importantly, that early-season behavior of other bird species makes them more available. Behaviors likely to influence avian host availability in the early season relate to reproduction, such as egg incubation and brooding of young. In Alabama, mosquito blood meals identified from herons peaked during the height of heron nesting season (Burkett-Cadena et al. 2011), whereas in Tennessee, the number of blood meals from robins oscillated up and down over the season, possibly in response to this species’ multiple broods per year (Savage et al. 2007). Nestlings have been considered especially susceptible to mosquitoes due to their decreased defensive behaviors, feathers, and motility; however, comparisons of landing and feeding rates on nestlings versus adults did not find evidence for this, possibly because adults shield them from bites during brooding (Griffing et al. 2007, Burkett-Cadena et al. 2010).

Instead, adult birds sitting on the nest for long periods of time may themselves become easy targets for mosquitoes. For example, Cooper’s Hawks have an incubation time of over a month during which the female is on the nest day and night, leaving only when fed periodically by the male (Baicich and Harrison 1997, Poole 2005). In New York, this typically

occurs from late April to early May (Meng 1951). Our samples collected during this time period showed a large number of Cooper's Hawk blood meals compared to other studies, the majority of which (8/10 sexed meals) are female. In addition, we found a significant bias toward female blood meals across all species in which the female alone performs incubation duties, suggesting that *Cx. restuans* targets incubating birds for blood feeding. This could explain why we observed a greater diversity of hosts in the early season, as the assortment of nest-bound birds in early spring could act like a "buffet for mosquitoes," whereas hosts are more difficult to locate and target once nestlings fledge and nests are abandoned.

The dilution effect hypothesis predicts that greater host diversity should result in lesser disease incidence (Ostfeld and Keesing 2000). This has been shown for WNV and nonpasserine diversity (Ezenwa et al. 2006, Allan et al. 2009) because nonpasserines tend to be poor WNV hosts (Komar et al. 2003, Blitvich 2008). Our dataset contains a large number of meals from Cooper's Hawks and Mourning Doves, which as Accipitriformes and Columbiformes, respectively, are poor hosts of WNV (Komar et al. 2003). We also detected blood meals from wetland birds (e.g., Great Egret, Ross's Goose), and areas with large colonies of wetland birds have been associated with lower mosquito infection rates (Reisen et al. 2009).

## Conclusions

Our study reveals that specific bird behaviors associated with the time of year, such as egg incubation in the spring, may be critical drivers of WNV amplification dynamics. This is in contrast to current hypotheses that WNV amplification hinges on the occurrence of specific vector species, such as *Cx. pipiens*. In fact, *Cx. restuans* is similar to *Cx. pipiens* in its late season host usage and therefore their different roles in WNV transmission may result from differences in seasonal timing rather than innate species characteristics.

## Acknowledgments

We thank Robert Fleischer (Smithsonian Institution) for information about sexing birds. We thank the following for sample collection and identification: Linda McCuiston, Rutgers University; Tony Acquaviva, Monmouth County Mosquito Extermination Commission; Peter Bosak and Diane McNelly, Cape May Mosquito Control; Christine Musa and Sara May, Warren County Mosquito Extermination Commission; Bill Reinert and Dana Chort, Atlantic County Mosquito Control; Rich Candeletti and Mike Romanowski, Ocean County Mosquito Extermination Commission; Sean Devaney, Union County Mosquito Control; and the Salem County Mosquito Extermination Commission. This work was funded by USDA Hatch Grant no. NJ08194, NE-1043 Multistate funds, and start-up funds to D.M. Fonseca.

## Author Disclosure Statement

No competing financial interests exist.

## References

Allan BF, Langerhans RB, Ryberg WA, Landesman WJ, et al. Ecological correlates of risk and incidence of West Nile virus in the United States. *Oecologia* 2009; 158:699–708.

- Andreadis TG, Anderson JF, Vossbrinck CR. Mosquito surveillance for West Nile virus in Connecticut, 2000: Isolation from *Culex pipiens*, *Cx. restuans*, *Cx. salinarius*, and *Culiseta melanura*. *Emerg Infect Dis* 2001; 7:670–674.
- Apperson CS, Harrison BA, Unnasch TR, Hassan HK, et al. Host-feeding habits of *Culex* and other mosquitoes (Diptera: Culicidae) in the Borough of Queens in New York City, with characters and techniques for identification of *Culex* mosquitoes. *J Med Entomol* 2002; 39:777–785.
- Apperson CS, Hassan HK, Harrison BA, Savage HM, et al. Host feeding patterns of established and potential mosquito vectors of West Nile virus in the eastern United States. *Vector Borne Zoonotic Dis* 2004; 4:71–82.
- Baichich PJ, Harrison CJO. *A Guide to the Nests, Eggs, and Nestlings of North American Birds*. California: Natural World Academic Press, 1997.
- Blitvich BJ. Transmission dynamics and changing epidemiology of West Nile virus. *Anim Health Res Rev* 2008; 9:71–86.
- Boyle WJ. *The Birds of New Jersey: Status and Distribution*. Princeton, NJ: Princeton University Press, 2011.
- Burkett-Cadena ND, Ligon RA, Liu M, Hassan HK, et al. Vector-host interactions in avian nests: Do mosquitoes prefer nestlings over adults? *Am J Trop Med Hyg* 2010; 83:395–399.
- Burkett-Cadena ND, McClure CJ, Ligon RA, Graham SP, et al. Host reproductive phenology drives seasonal patterns of host use in mosquitoes. *PLoS One* 2011; 6:e17681.
- Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West Nile virus. *Lancet Infect Dis* 2002; 2:519–529.
- Cicero C, Johnson NK. Higher-level phylogeny of new world vireos (Aves: Vireonidae) based on sequences of multiple mitochondrial DNA genes. *Mol Phylogenet Evol* 2001; 20:27–40.
- Crabtree MB, Savage HM, Miller BR. Development of a species-diagnostic polymerase chain reaction assay for the identification of *Culex* vectors of St. Louis encephalitis virus based on interspecies sequence variation in ribosomal dna spacers. *Am J Trop Med Hyg* 1995; 53:105–109.
- Cupp EW, Zhang D, Yue X, Cupp MS, et al. Identification of reptilian and amphibian blood meals from mosquitoes in an eastern equine encephalomyelitis virus focus in central Alabama. *Am J Trop Med Hyg* 2004; 71:272–276.
- Diuk-Wasser MA, Molaei G, Simpson JE, Folsom-O'Keefe CM, et al. Avian communal roosts as amplification foci for West Nile virus in urban areas in northeastern United States. *Am J Trop Med Hyg* 2010; 82:337–343.
- Dohm DJ, O'Guinn ML, Turell MJ. Effect of environmental temperature on the ability of *Culex pipiens* (Diptera: Culicidae) to transmit West Nile virus. *J Med Entomol* 2002; 39:221–225.
- Dunne P, Kane R, Kerlinger P. *New Jersey at the Crossroads of Migration*. Cape May Point, NJ: New Jersey Audubon Society, 1989.
- Ebel GD, Rochlin I, Longacker J, Kramer LD. *Culex restuans* (Diptera: Culicidae) relative abundance and vector competence for West Nile virus. *J Med Entomol* 2005; 42:838–843.
- Egizi A, Healy SP, Fonseca DM. Rapid blood meal scoring in anthropophilic *Aedes albopictus* and application of PCR blocking to avoid pseudogenes. *Infect Genet Evol* 2013; 16:122–128.
- Ezenwa VO, Godsey MS, King RJ, Guptill SC. Avian diversity and West Nile virus: Testing associations between biodiversity and infectious disease risk. *Proc Biol Sci* 2006; 273:109–117.
- Gong H, DeGaetano A, Harrington L. Climate-based models for West Nile *Culex* mosquito vectors in the Northeastern US. *Int J Biometeorol* 2011; 55:435–446.

- Griffing SM, Kilpatrick AM, Clark L, Marra PP. Mosquito landing rates on nesting American robins (*Turdus migratorius*). *Vector Borne Zoonotic Dis* 2007; 7:437–443.
- Griffiths R, Double MC, Orr K, Dawson RJG. A DNA test to sex most birds. *Mol Ecol* 1998; 7:1071–1075.
- Hamer GL, Kitron UD, Brawn JD, Loss SR, et al. *Culex pipiens* (Diptera: Culicidae): A bridge vector of West Nile virus to humans. *J Med Entomol* 2008; 45:125–128.
- Hamer GL, Kitron UD, Goldberg TL, Brawn JD, et al. Host selection by *Culex pipiens* mosquitoes and West Nile virus amplification. *Am J Trop Med Hyg* 2009; 80:268–278.
- Johnson B, Munafo K, Shappell L, Tsipoura N, Robson M, Ehrenfeld J, Sukhdeo MK. The roles of mosquito and bird communities on the prevalence of West Nile virus in urban wetland and residential habitats. *Urban Ecosystems* 2012; 15:513–531.
- Kent R, Juliusson L, Weissmann M, Evans S, et al. Seasonal blood-feeding behavior of *Culex tarsalis* (Diptera: Culicidae) in Weld County, Colorado, 2007. *J Med Entomol* 2009; 46:380–390.
- Kent RJ. Molecular methods for arthropod bloodmeal identification and applications to ecological and vector-borne disease studies. *Mol Ecol Resour* 2009; 9:4–18.
- Kilpatrick AM, Daszak P, Jones MJ, Marra PP, et al. Host heterogeneity dominates West Nile virus transmission. *Proc R Soc B Biol Sci* 2006a; 273:2327–2333.
- Kilpatrick AM, Kramer LD, Jones MJ, Marra PP, et al. West Nile virus epidemics in North America are driven by shifts in mosquito feeding behavior. *PLoS Biol* 2006b; 4:e82.
- Kilpatrick AM, Meola MA, Moudy RM, Kramer LD. Temperature, viral genetics, and the transmission of West Nile virus by *Culex pipiens* mosquitoes. *PLoS Pathog* 2008; 4:e1000092.
- Kindt R, Coe R. Tree diversity analysis. *A Manual and Software for Common Statistical Methods for Ecological and Biodiversity Studies*. Nairobi: World Agroforestry Centre (ICRAF), 2005.
- Kocher TD, Thomas WK, Meyer A, Edwards SV, et al. Dynamics of mitochondrial DNA evolution in animals: Amplification and sequencing with conserved primers. *Proc Natl Acad Sci USA* 1989; 86:6196–6200.
- Komar N, Langevin S, Hinten S, Nemeth N, et al. Experimental infection of North American birds with the New York 1999 strain of West Nile virus. *Emerg Infect Dis* 2003; 9:311–322.
- Kramer LD, Styer LM, Ebel GD. A global perspective on the epidemiology of West Nile virus. *Annu Rev Entomol* 2008; 53:61–81.
- Lampman R, Slamecka M, Krasavin N, Kunkel K, et al. *Culex* population dynamics and West Nile Virus transmission in East-Central Illinois. *J Am Mosq Control Assoc* 2006; 22:390–400.
- Lee JH, Rowley WA. The abundance and seasonal distribution of *Culex* mosquitoes in Iowa during 1995–97. *J Am Mosq Control Assoc* 2000; 16:275–278.
- Mabey SE, McCann JM, Niles LJ, Barlett C, Kerlinger P. Neotropical Migratory Songbird Regional Coastal Corridor Study Final Report. Virginia Department of Environmental Quality, 1993.
- Meng RK. *The Cooper's hawk, Accipiter cooperii (Bonaparte)*. Ithaca, NY: Cornell University, 1951:202.
- Molaei G, Andreadis TG, Armstrong PM, Anderson JF, et al. Host feeding patterns of *Culex* mosquitoes and West Nile virus transmission, northeastern United States. *Emerg Infect Dis* 2006; 12:468–474.
- Montgomery MJ, Thiemann T, Macedo P, Brown DA, et al. Blood-feeding patterns of the *Culex pipiens* complex in Sacramento and Yolo Counties, California. *J Med Entomol* 2011; 48:398–404.
- Nasci RS, Komar N, Marfin AA, Ludwig GV, et al. Detection of West Nile virus-infected mosquitoes and seropositive juvenile birds in the vicinity of virus-positive dead birds. *Am J Trop Med Hyg* 2002; 67:492–496.
- Nenadic O, Greenacre M. Correspondence Analysis in R, with two- and three-dimensional graphics: The ca package. *J Statist Software* 2007; 20:1–13.
- Ngo KA, Kramer LD. Identification of mosquito bloodmeals using polymerase chain reaction (PCR) with order-specific primers. *J Med Entomol* 2003; 40:215–222.
- Oksanen J, Blanchet FG, Kindt R, Legendre P, et al. *vegan: Community Ecology Package*. R package version 2.0-7, 2013.
- Ostfeld RS, Keesing F. Biodiversity and disease risk: The case of Lyme disease. *Conserv Biol* 2000; 14:722–728.
- Patrican LA, Hackett LE, Briggs JE, McGowan JW, et al. Host-feeding patterns of *Culex* mosquitoes in relation to trap habitat. *Emerg Infect Dis* 2007; 13:1921–1923.
- Pielou E. The measurement of diversity in different types of biological collections. *J Theoret Biol* 1966; 13:131–144.
- Poole A. *The Birds of North America Online*: <http://bna.birds.cornell.edu/BNA/>. Ithaca, NY: Cornell Laboratory of Ornithology, 2005.
- R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing, 2013.
- Reisen WK, Wheeler S, Armijos MV, Fang Y, et al. Role of communally nesting ardeid birds in the epidemiology of West Nile virus revisited. *Vector Borne Zoonotic Dis* 2009; 9:275–280.
- Reiter P. Weather, vector biology, and arboviral recrudescence. In: Monath TP, ed. *Arboviruses: Epidemiology and Ecology*. Boca Raton, FL: CRC Press, 1988:245–255.
- Rényi A. On measures of information and entropy. *Proceedings of the fourth Berkeley Symposium on Mathematics, Statistics and Probability* 1960 1961:547–561.
- Savage HM, Aggarwal D, Apperson CS, Katholi CR, et al. Host choice and West Nile virus infection rates in blood-fed mosquitoes, including members of the *Culex pipiens* complex, from Memphis and Shelby County, Tennessee, 2002–2003. *Vector Borne Zoonotic Dis* 2007; 7:365–386.
- Shannon CE, Weaver W. *The Mathematical Theory of Communication*. Urbana: University of Illinois Press; 1949.
- Simpson EH. Measurement of species diversity. *Nature* 1949; 163:688.
- Simpson JE, Folsom-O'Keefe CM, Childs JE, Simons LE, Andreadis TG, Diuk-Wasser MA. Avian host-selection by *Culex pipiens* in experimental trials. *PLoS One* 2009; 4:e7861.
- Tóthmérész B. Comparison of different methods for diversity ordering. *J Vegetat Sci* 1995; 6:283–290.

Address correspondence to:  
 Dina M. Fonseca  
 Center for Vector Biology  
 Rutgers University  
 180 Jones Avenue  
 New Brunswick, NJ 08901

E-mail: dinafons@rci.rutgers.edu